

Goals

- Direct sum
- Orthogonal complement
- Orthogonal projection

Defn For U, V F -vector spaces, their direct sum is

\oplus

$$U \oplus V := U \times V = \{(u, v) \mid u \in U, v \in V\}$$

with componentwise addition and scalar multiplication:

$$(u, v) + (u', v') = (u + u', v + v'),$$

$$\lambda(u, v) = (\lambda u, \lambda v).$$

E.g. $\mathbb{R}^2 = \mathbb{R} \oplus \mathbb{R}$

Prop Let $U, V \leq W$ such that

① $\text{span}(U \cup V) = W$,

② $U \cap V = \{0\}$.

Then $U \oplus V \xrightarrow{\cong} W$

$(u, v) \mapsto u + v$

Write $W = U \oplus V$

PF Linear ✓

① guarantees surjectivity. If $(u, v) \in \ker$ then $u + v = 0 \Rightarrow u = -v \in U \cap V \Rightarrow u = v = 0$ by ②.

Thus $\ker = \{0\}$ so the map is injective as well. $\square$


From now on, $(V, \langle \cdot, \cdot \rangle)$ an inner product space over $F = \mathbb{R}$ or $\mathbb{C}$.

Defn For $S \subseteq V$, the orthogonal complement of S is

$$S^\perp := \{x \in V \mid \langle x, s \rangle = 0 \ \forall s \in S\}$$

Q Is $S^\perp$ a subspace of V ?

$S = \{s\}$ 



A $0 \in S^\perp$ ✓. If $x, y \in S^\perp$, $\langle x + \lambda y, s \rangle = \langle x, s \rangle + \lambda \langle y, s \rangle$
 $= 0 + \lambda 0 = 0 \quad \forall s \in S \Rightarrow x + \lambda y \in S^\perp$.

Prop Suppose $\dim V = n$, $S = \{v_1, \dots, v_k\} \subseteq V$ is orthonormal.

① S can be extended to an orthonormal basis $\{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$ of V .

② If $W = \text{span } S$, then $\{v_{k+1}, \dots, v_n\}$ is an orthonormal basis of $S^\perp = W^\perp$.

③ If $W \subseteq V$, then $\dim W + \dim W^\perp = \dim V = n$. $y \in x$

④ If $W \subseteq V$, then $(W^\perp)^\perp = W$. (Similar to $x \cdot (x - y) = y$.)

Pf ① Apply G-S to any basis extension of $\{v_1, \dots, v_k\}$.

② Let $S' = \{v_{k+1}, \dots, v_n\}$, which is lin ind & orthonormal. Since S is orthonormal, $S' \subseteq W^\perp \Rightarrow \text{span } S' \subseteq W^\perp$. For $x \in W^\perp$, since $S \cup S'$ is orthonormal,

$$\begin{aligned}
 x &= \sum_{i=1}^n \langle x, v_i \rangle v_i \\
 &= \sum_{i=k+1}^n \langle x, v_i \rangle v_i \quad [x \in W^\perp] \\
 &\in \text{span } S'
 \end{aligned}$$

so $\text{span } S' = W^\perp \Rightarrow S'$ is a basis for $W^\perp$.

③ Follows directly from ②.

④ We have $(W^\perp)^\perp = \{x \in V \mid \langle x, y \rangle = 0 \text{ for } y \in W^\perp\} \supseteq W$.

Now $\dim(W^\perp)^\perp \stackrel{\textcircled{3}}{=} n - \dim W^\perp \stackrel{\textcircled{3}}{=} \dim W$ so $W = (W^\perp)^\perp$. $\square$

Prop Let $W \leq V$. Then $V = W \oplus W^\perp$, i.e. $\perp$ for all $y \in V$ there is a unique $u \in W, v \in W^\perp$ such that $y = u + v$.

Call u the orthogonal projection of y onto W . If $\{u_1, \dots, u_k\}$ is an

orthonormal basis for W , then $u = \sum_{i=1}^k \langle y, u_i \rangle u_i$.

Pf Let $\{u_1, \dots, u_k\}$ be an orthonormal basis for W and define $u = \sum_{i=1}^k \langle y, u_i \rangle u_i$, $v = y - u$. Then $u \in W$ and $y = u + v$.

Furthermore, for $1 \leq j \leq k$,

$$\begin{aligned} \langle v, u_j \rangle &= \langle y - u, u_j \rangle \\ &= \langle y, u_j \rangle - \left\langle \sum_{i=1}^k \langle y, u_i \rangle u_i, u_j \right\rangle \\ &= \langle y, u_j \rangle - \sum_{i=1}^k \langle y, u_i \rangle \langle u_i, u_j \rangle \\ &= \langle y, u_j \rangle - \langle y, u_j \rangle \cdot 1 \quad [\text{orthonormality}] \\ &= 0 \end{aligned}$$

so $v \in W^\perp$. Moral exercise: check $W \cap W^\perp = \{0\}$ and $\checkmark$ thus the expression is unique. $\square$


Cor The orthogonal projection u of y onto W is the vector in W closest to y : $\|y-u\| \leq \|y-w\|$ for all $w \in W$ with equality iff $u=w$.

PF Write $y = u + v$ with $u \in W$, $v \in W^\perp$. Take $w \in W$. Then $u-w \in W$, $y-u \in W^\perp$ so by Pythagoras,

$$\|y-w\|^2 = \|y-u + u-w\|^2 \quad [\text{Get Smart}]$$

$$= \|y-u\|^2 + \|u-w\|^2 \quad [\text{Pythag}]$$

$$\geq \|y-u\|^2 \quad [\|u-w\|^2 \geq 0]$$




with equality iff $\|u-w\| = 0$ iff $u=w$. $\square$

E.g. Let $W = \text{span}\{ \overset{u_1}{(1,1,0)}, \overset{u_2}{(0,0,1)} \}$. Determine the distance of $y = (4,0,-1)$ from W :

$$u = \frac{\langle y, u_1 \rangle}{\|u_1\|^2} u_1 + \frac{\langle y, u_2 \rangle}{\|u_2\|^2} u_2 = 2\sqrt{2} u_1 + \frac{-\sqrt{2}}{2} u_2$$

$$= (2\sqrt{2}, 2\sqrt{2}, -\frac{\sqrt{2}}{2})$$

$$\text{so } y-u = (4-2\sqrt{2}, -2\sqrt{2}, -1+\frac{\sqrt{2}}{2}) \text{ with } \|y-u\| = \sqrt{\frac{67}{2} - 17\sqrt{2}} \approx 3.075$$



Topics

- Spectral theorem
- Markov chains $\rightarrow$ Page Rank
- cross product
- matrix groups $SO(n)$ (esp topology $SO(3)$)
 $SL_2\mathbb{R}$
- normed division algebras : quaternions
- SVD - singular value decomposition
- your ideas!!